

University of Groningen

Hansenula polymorpha pex11 cells are affected in peroxisome retention

Krikken, Arjen M; Veenhuis, Marten; van der Klei, Ida J

Published in:
Febs Journal

DOI:
[10.1111/j.1742-4658.2009.06883.x](https://doi.org/10.1111/j.1742-4658.2009.06883.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Krikken, A. M., Veenhuis, M., & van der Klei, I. J. (2009). Hansenula polymorpha pex11 cells are affected in peroxisome retention. *Febs Journal*, 276(5), 1429-1439. <https://doi.org/10.1111/j.1742-4658.2009.06883.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary material

Video 1. Time-lapse imaging of peroxisomes in *H. polymorpha* wild type cells growing in the presence of glucose. Cells produce GFP-SKL to visualize peroxisomes. Peroxisomes are equally distributed over mother and daughter cell at an early stage of bud development. The video was captured at 30 frames/hour and is displayed at 2 frames/second. Video stills from this video are presented in Figure 3A.

Video 2. Time-lapse imaging of peroxisomes in *H. polymorpha pex11* cells growing in the presence of glucose. Cells produce GFP-SKL to visualize peroxisomes. The peroxisome migrates into the developing bud at an early stage of bud development leaving the mother cell devoid of peroxisomes. The video was captured at 30 frames/hour and is displayed at 2 frames/second. Video stills from this video are presented in Figure 3B.

Video 3. Time-lapse imaging of peroxisomes in *H. polymorpha* wild type cells producing Inp1-GFP growing in the presence of glucose. Upon initiation of bud formation the Inp1-GFP fluorescence increased strongly to the organelles that are retained in the mother cells. The video was captured at 20 frames/hour and is displayed at 2 frames/second. Video stills from this video are presented in Figure 5A.

Video 4. Time-lapse imaging of *H. polymorpha pex11* cells producing Inp1-GFP growing in the presence of glucose. In *pex11* cells Inp1-GFP is localized to peroxisomes that are transported to the bud. The video was captured at 20 frames/hour and is displayed at 2 frames/second. Video stills from this video are presented in Figure 5B.